

Artisanal and Small-scale Gold Mining in Syanyonja, Busia Gold District, South Eastern Uganda: Impacts on the Mining Population and the Environment

Timothy Omara^{1,2*}, Shakilah Karungi³, Stephen Ssebulime⁴,
Kibet Mohamed Kiplagat⁵, Ocident Bongomin⁶, Remish Ogwang⁷
and Solomon Akaganyira⁸

¹Department of Chemistry and Biochemistry, School of Biological and Physical Sciences, Moi University, Uasin Gishu County, Kesses, P.O.Box 3900-30100, Eldoret, Kenya.

²Department of Quality Control and Quality Assurance, Product Development Directory, AgroWays Uganda Limited, Kyabazinga Way, P.O.Box 1924, Jinja, Uganda.

³Department of Mining and Water Resources Engineering, Faculty of Engineering, Busitema University, P.O.Box 236, Busitema, Tororo, Uganda.

⁴Directorate of Government Analytical Laboratory, Ministry of Internal Affairs, Lourdel Rd, Wandegeya, P.O.Box 2174, Kampala, Uganda.

⁵Department of Mechanical Engineering, School of Engineering, Moi University, Uasin Gishu County, Kesses, P.O.Box 3900-30100, Eldoret, Kenya.

⁶Department of Manufacturing, Industrial and Textile Engineering, School of Engineering, Moi University, Uasin Gishu County, Kesses, P.O.Box 3900-30100, Eldoret, Kenya.

⁷Department of Quality Control, Rene Industries Limited, P.O.Box 6034, Kampala, Uganda.

⁸Department of Mechanical Engineering, Faculty of Engineering and Architecture, American University of Beirut, P.O.Box 11-0236, Riad El-Solh 1107 2020, Beirut, Lebanon.

Authors' contributions

This work was carried out in collaboration among all authors. Authors TO and SK designed the study and performed fieldwork. Authors SS and OB organized data and performed the statistical analysis. Authors TO, SK, KMK, OB, RO and SA wrote the first draft of the manuscript. Authors TO, KMK and OB managed the analyses of the study. Authors TO, SK, SS and SA managed the literature searches. All authors read and approved the final manuscript.

Article Information

Editor(s):

(1) Dr. Adewumi, Adeniyi JohnPaul, Lecturer II, Department of Geological Sciences, Achievers University, Owo, Nigeria.

Reviewers:

(1) Antipas T. S. Massawe, University of Dar es Salaam, Tanzania.

(2) Gwisai Reginald Dennis, Bindura University of Science Education, Zimbabwe.

(3) Acaye Genesis, Cyan International, India.

Complete Peer review History: <https://sdiarticle4.com/review-history/52582>

Original Research Article

Received 01 September 2019
Accepted 03 November 2019
Published 11 November 2019

ABSTRACT

Aims: To determine the amount of mercury discharged into Namukombe stream, the major water body in Syanyonja village, Busia gold district and investigate the impacts of mercury-based artisanal and small-scale gold mining (ASGM) on the mining population and the environment.

Study Design: Quantitation of mercury discharged with tailings was done by mass balance method. Field survey at the mining sites was done followed by administration of questionnaires to 50 stamperders in the village.

Place and Duration of Study: The study was done in Syanyonja village in Syanyonja parish, Busitema subcounty, Samia Bugwe constituency, Busia gold district, South East of the Republic of Uganda between April 2019 to June 2019.

Methodology: Fieldwork was done in Syanyonja village to appreciate the level of environmental pollution due to mercurial ASGM in the area. Quantitation of mercury discharged with tailings was done by mass balance method. Field survey at the mining sites was done followed by administration of 50 questionnaires to stamperders of at least 18 years old and ASGM experience of not less one year.

Results: About 8% of mercury mixed with auriferous materials is lost in tailings, accounting for an annual mercury release of about 1.757 kg into the environment. Socio-demographic results indicated that the majority of the mining population (64%) are male and ASGM have left human health and environmental footprints, which directly or indirectly affects the population. The most common health problems among miners are malaria (36%) and abdominal pain (20%). The standard of living of the miners are evidently low, and most mines are characterized by school dropouts, prostitutes and thieves. Mining sites have deplorably poor hygiene, with evident burning of amalgams to recover gold.

Conclusion: ASGM have been accompanied by wanton mowing down of vegetation, land degradation as well as mercuric pollution of water, air, land and aquatic ecosystems. It is suggested that the Ugandan government should re-enforce committees to follow up on ASGM activities and train artisans on sustainable (non-mercury) alternative gold extraction methods that do not create new toxic exposures such as using borax, magnets and sluice boxes. Artisans should take up farming actively as an alternative.

Keywords: Artisanal gold mining; Busia; environmental impacts; mercury pollution; Namukombe stream; panning; Syanyonja; Uganda.

1. INTRODUCTION

Uganda is a low-income landlocked African country with an averagely solid economic growth and demonstrated ambitions to attain a middle-income status by 2020 [1]. The country is at a juncture in its history of economic transformation preceding a decade of an apogee of Gross Domestic Product (GDP) growth that calamitously dwindled from 7% in 2009 to 3.9% in the fiscal year 2016/2017 [2,3]. Accordingly, the fiscal policy is badly advocating for capital investment to revitalize this GDP. The minerals sector which used to be informal and largely illegal is poised to contribute a pivotal part in this economic move as evidenced by its position of citation in the second National Development Plan for 2015 to 2020 [1].

Uganda lies in the African plate, a continental crust rich in Precambrian rocks of Achaean and Lower Proterozoic to Middle Proterozoic geology

[4]. Thus, it has a substantial geological potential for precious minerals including copper, cobalt, tungsten, beryllium, bismuth, uranium, limestone, phosphate, vermiculite, kaolin, bentonite, diatomite, gypsum, glass sands, salts, pozzolana, dimension stones, gemstones, gold and crude oil [1,4-6]. Artisanal and small scale gold mining (ASGM) has been reported in Ugandan gold districts of Mubende (Kamalenge and Kisita), Kabale (Chiruruma, Chilima, Bugarama, Mugyera), Kisoro (Murindi, Mpororo, Rubuguri and Karamba), Namayingo, Bugiri, Mayuge, Busia (Tiira, Syanyonja, Makina, Amonikakinei), Buhweju (Kahengyere and Muti), Rukungiri (Bikongozo valley), Bushenyi (Mashonga in Kyamuhunga), Mbarara (Kitaka), Ibanda, Kanungu (Kashenyi, Kanungu, and Muramba), Moroto (Rupa, Katikikele and Kamalera), Abim (Alarek and Morulem), Nakapiripirit (Acherer), Kaabong (Lopedo, Sokodo, Kanguoth, Lemoyne, Nakapiril-Kekuli and Lugoto), Nakapiripirit, Hoima (Kafu River)

and Amudat (Karita) [4,7,8]. Except in Kisita, Kamalenge, Tiira and Amonikakinei where gold is recovered from reefs, gold is often recovered from alluvial auriferous materials, with Busia being the pioneer gold mining district [4,6].

The Ugandan government recently accorded artisanal mining greater cognizance through its prominent inclusion in a draft of the National Mining and Minerals Policy. This initiative is believed shall establish the basis for the upcoming mining laws and regulations and related strategies and actions by the government. In view of the foregoing, more robust evidence of the significance of artisanal mining and clarity on the opportunities and challenges presented by the sector are needed.

Busia have underground workings and surface mining of both alluvial and hard rock gold ores. The principal livelihood activities in this area include trade (as it is at the border with Kenya), gold mining and agriculture [9]. Up to 600 vulnerable people of Buteba, Sikuda and Busitema sub-counties in Busia subsist on ASGM. A demographic ratio of 4:1 (men to women) was reported, with individual annual earnings estimated at 17 million Uganda shillings [10].

Globally, the upsurge of ASGM and the growing awareness of the negative impacts of mercury use, both in terms of environmental and human health, has culminated in the drafting and ratification of the Minamata Convention which constrain signatories to phase out mercury use. As a contribution to decision making, this study evaluated the impacts of mercury-based ASGM in Uganda, taking evidence from Namukombe stream and its propinquity (Syanyonja village) as an example due to the intermittent ASGM in this area.

2. MATERIALS AND METHODS

2.1 Brief Description of the Study Area

The study was conducted in Syanyonja village, Syanyonja Parish, Busitema subcounty, Samia Bugwe constituency, Busia gold district, South East of the Republic of Uganda. The choice of this area is explained by the high intensity of mercurial ASGM in it. Busia lies between 33°05' E 00°10' N and 34°01'E 00°35'N [6], covering an expanse of about 730.9 km² of land. It is flanked by Tororo district to the North, Kenya to the East, Lake Victoria to the South, Namayingo district to the Southwest and Bugiri district to the West [6].

Busia comprises of a mixture of auriferous hard rock (70%) and 30% alluvial soil which are part of the Cretaceous to Miocene intrusive alkaline carbonatite complexes of the Ugandan frontier with Kenya [4]. Namukombe stream, a tributary of River Malaba is one of the water bodies in Syanyonja village characterized by mercury-based ASGM activities [6] and was of interest in this study.

2.2 Apparatus Used

A calibrated Mettler PM200 analytical balance (Marshall Scientific, Hampton, USA), retort, single use disposable hand gloves, dust masks and eye googles.

2.3 Determination of Amount of Mercury Discharged with Slurry Tailings

Informal interviews with artisanal gold miners (stampedeers) was conducted to assess the mercury losses during gold recovery. Mercury recovered after squeezing and panning the amalgams were estimated by mercury mass balance method described by preceding investigations [11,12].

The mass of mercury (quicksilver) entering the system (X g) was measured using an analytical balance before being introduced into the amalgamation process to concentrate gold with heavier particles in a pan. The mass of the solid mercury-gold amalgam formed (Y g) and the amount of pure gold recovered (Z g) were measured after amalgamation and panning respectively. The mass of mercury recovered after amalgamation (excess mercury), M g was determined from the mass of mercury recovered after squeezing the amalgam. The amalgam was burnt in open air on a retort to evaporate the mercury and then the solid was re-weighed and the mass noted.

The mass of mercury lost in the slurry tailings (discharged), D g was estimated from the difference in the masses of the mercury measured utilizing Equation 1.

$$D = X - (Z - Y) - M \quad (1)$$

The percentage of mercury discharged in slurry tailings was calculated from Equation 2.

$$\% \text{ discharged} = \frac{D}{X} \times 100\% \quad (2)$$

The assumptions made were that no mercury was lost through evaporation during



Fig. 1. Administration of questionnaires in Syanyonja village

amalgamation and losses due to adsorption onto pan walls are negligible. The procedure was repeated using different masses (2, 3, 4, 5 and 6 g) of mercury which are typical of the quantities of mercury used for amalgamation at different ASGM sites in Syanyonja village.

2.4 Collection of Field Data

Direct field observation was done in the mining sites in order to appreciate the rate of environmental destruction and the various phases involved in mercurial ASGM. The fieldwork in Syanyonja village was conducted between April 2019 and June 2019. Stampeders (at least 18 years old with a seniority of at least 1 year and consenting) were selected. Because of the absence of an exhaustive list of stampeders working on the ASGM sites due to the intermittent nature of this activity, selection by simple random sampling was impossible. A quota sampling according to sex (1/3 of women, 2/3 of men) was performed considering the fact that approximately one-third of women work in ASGM sites [13]. To compensate for the selection bias associated with this type of sampling, the estimated size of the sample ($n = 25$) was doubled to 50 respondents to cover potential non-respondents. In quantitative terms, the power of the target study was at least 75%, sufficient in the event where less than 50% of the stampeders are unaware of the impacts of mercury use in ASGM on miners and the environment [14]. Individual interviews (face to

face) were conducted basing on an anonymous, standardized questionnaire (Fig. 1). The questionnaire captured socio-demographic characteristics, common illness in the area, number of family members engaged in ASGM, types and sources of proteinaceous food eaten and the sources of water for domestic use (drinking) and for livestock.

2.5 Statistical Analysis of Results

All data collected in this study were entered into Microsoft Excel for Office 365 (version 16, Microsoft Corporation, USA). Statistical analyses (One-way ANOVA, averaging and other descriptive statistics) were done using Minitab statistical software (Version 17, Minitab Inc., USA). Statistical significance was set at $P = .05$.

3. RESULTS AND DISCUSSION

3.1 Mercury Lost in Slurry Tailings

The results of mercury discharged with tailings are given in Table 1.

About 8% of mercury used in amalgamation is discharged into the environment through slurry tailings during gold processing in Syanyonja and this poses a great threat to the environment. According to the interviewed mercury sellers around the mining sites, an average of 60 g of mercury is sold and used daily, implying that 4.8 g and 1,756.8 g of mercury are discharged into

the slurry tailings daily and annually respectively. A report [13] indicated that about 58% ($n = 118$) of sampled gold stampeder in a study done in Mubende and Busia gold districts of Uganda used about 1–50 g of mercury per person weekly. This computationally translates into 2,800 g per stampeder per year. The results of this study is in complete agreement with the anecdotal finding of COWI [15] which observed that, although there were no recent estimates on the use of mercury for ASGM in Uganda, the mean mercury consumption per miner from Busia and Mubende gold-mining sites could give a mercury consumption rate of about 40.5 tonnes per year. This is further substantiated by an estimate which indicated that approximately 1,400 tons of mercury are used every year by ASGM miners globally [16]. In Rwamagasa ASGM area located in Geita district of Tanzania, about 27 kg of mercury was estimated to be released to the environment yearly, with about 12 kg of gold recovered from the process [17]. It is equally likely that the amount of mercury lost to the slurry tailings is re-emitted to the atmosphere on a relatively short time scale of one to several years as the tailings are accumulated in above-the-ground piles and later reprocessed. It should therefore be over emphasized that the population living near to or downstream of Namukombe stream and the ASGM sites are at risk of mercury pollution [6].

3.2 Socio-demographic Characteristics and Exposure to Mercury

A total of 50 stampeder were contacted in this study. The characteristics of the participants are given in Table 2. The average age of the respondents was 26 years. The sample consisted of 64% men ($n = 32$) and 36% women ($n = 18$). There was no significant difference ($P = .05$) in the number of people who had stayed for 1 to 5 years and over 16 years in Syanyonja village (Table 2). Most of the family members of

the respondents (80%) were either directly or indirectly involved in mercury based ASGM. The gold mines were dominated by people between 20 years and above (70%) though children (stampeder below 18 years of age) constituted about 14% of the mining population (Table 2). Overall, the levels of child labour were higher in the ASGM sites and were more informal where participation were poverty rather than enterprise driven. Some of the children in these mines are orphans, are from child-led households or lack school fees and scholastic materials, corroborating preceding studies in other parts of Uganda [7,18]. This study also agreed well with previous reports that women realize fewer benefits from artisanal and small scale mining than men but bear the brunt of negative impacts [7,19-22].

In addition, the mines consisted of people of various ethnic origins namely: Bagishu, Basoga, the Iteso, Karamojong (Karimojong) and the Congolese. This is similar to a report [7] which cited that Acherer, an ASGM site in Karamoja region (Nakapiririt district) hosts different ethnic groups: the Iteso, Western Ugandans and Rwandese during peak periods of ASGM. It is known that decentralization, economic turmoil, the ten-fold gold price increase within the period of 1996 to 2002 coupled by prospects of improved livelihoods from ASGM attracts unemployed people to engage in ASGM [7,20, 23].

In terms of exposure, the findings indicated that most respondents (50%, $n = 25$) had spent 31 years or more using mercury, most of which (68%, $n = 17$) were women. In a similar cross-sectional study in some sites of Mubende and Busia districts of Uganda, it was reported that 44% ($n = 90$) of the respondents had used mercury for ASGM for 1 to 4 years while the remaining percentage (56%, $n = 114$) had lived for 5 to 29 years handling ASGM mercury [13].

Table 1. Mass balance for mercury lost in slurry tailings

Mercury added, X (g)	Mass of amalgam formed, Y (g)	Mass of gold recovered, Z (g)	Mass of mercury recovered after amalgamation, M(g)	Mass of mercury lost in tailings, D (g)
1	0.36	0.13	0.51	0.26
2	1.24	0.22	0.70	0.28
3	2.34	0.30	0.69	0.27
4	3.45	0.36	0.63	0.28
5	4.39	0.40	0.71	0.30
6	5.51	0.42	0.62	0.29
3.5±1.87*	2.88 ±1.94	0.31 ±0.11	0.64 ±0.08	0.28 ±0.14

*Results in this row are means of the corresponding columns with standard deviations attached

Table 2. Characteristics of the sampled respondents (n = 50)

Variable	Categories	Number	Percentage
Sex	Male	32	64
	Female	18	36
Residence time in Syanyonja village	1-5 years	22	45
	6-10 years	04	7
	11-15 years	02	3
	Over 16 years	22	45
Family members engaged in ASGM	Yes	40	80
	No	10	20
Age of stampeders engaged in ASGM*	1-10 years	04	8
	10-15 years	03	6
	15-20 years	08	16
	20-30 years	10	20
	31 years and over	25	50
Major sicknesses in the last 2 years	None	16	32
	Malaria	18	36
	Acute respiratory problems	02	4
	Abdominal pain	10	20
	Other health problems	04	8

*Ages cited were from averaged information provided by the respondents in regard to their family members engaged in ASGM

Pertaining health, it was noted that malaria (36%) and abdominal pain (20%) tops the list of the illnesses reported in the last two (2) years (Table 2). This could be because pits left after harvesting auriferous rock materials holds stagnant water which act as breeding grounds for female *anopheles*' mosquitoes, the *plasmodium* vector which enhances the spread of malaria-causing *plasmodium* (*P. falciparum* and *P. malariae*). Abdominal pains are probably due to the manual nature of ASGM in Syanyonja which involves digging, crushing and panning of auriferous materials, usually involving bending

for long hours in the scorching sun [6]. Gold artisans in Syanyonja employ rudimentary methods of gold extraction including panning [6], typical of other ASGM sites in Uganda [18,20, 21] (Fig. 2). Human health problems, diseases and neuropsychiatric disorders such as malaria, cholera, venereal diseases (HIV/AIDS, gonorrhoea, syphilis), musculoskeletal disorders (myalgia of the upper and lower limbs and the back), sleep disruption (insomnia), headache, anorexia, tremor of the fingers, loss of fecundity and abnormal fatigue have been reported in other ASGM communities [7,24-26].



Fig. 2. Panning of alluvium along Namukombe stream : (a) by children (school dropouts). Notice the plastic tin used for storing recovered gold nuggets; (b) by a man. Notice the ditches created; (c) quality of water from the stream. Notice the destruction of shrubs in the right background (modified from [6], photos reproduced with permission)

It was noted that in Syanyonja gold mines, women are multitasked. They actively engage in ore processing and transportation, water collection, catering, prostitution, operate clinics amongst others. Unfortunately, this pivotal role played by women is not recognized in official ASGM policies [27] and in some cases they end up being victims of rape [18]. Also, ASGM is combined with domestic and reproductive work as well as small scale farming. Most women and girls are involved in low paying jobs, as most high paying opportunities are usually work prohibited by cultural norms for females to do. Therefore, women usually end up in panning, collecting water, transporting auriferous rock materials for crushing and burning of mercury in open air. Some operate eateries, sell airtime and consumables (such as waragi, cigarettes, cakes) around the mines. This is corroborant with preceding studies in Uganda [7,18,21,27].

Children taken to the mining sites may be exposed to atmospheric mercury vapour resulting from burning of amalgams in open air. Inhalation of mercury vapour results in high blood concentration of mercury implicated for many respiratory, neurologic, renal, immunologic, dermatologic and reproductive complications [28]. Human health impacts emanating from the utilization of mercury in ASGM have been reported in African countries of Benin [29], Burkina Faso [30], Ghana [31,32] and Mali [33]. This is evidenced by elevated levels of mercury in human blood, hair, urine and breast milk [26,34,35].

ASGM is largely a poverty-driven activity, used as a coping mechanism with thousands of families engaging in the mining practice [18,36]. This unfortunately exposes the miners to dermal contact with mercury in the dredged sediments as well as mercury-contaminated water [6]. Like reported for other parts of Busia (Tiira and Amonikakinei) [37], Karamoja region [7, 18], Mubende and Buhweju [13,20,21], miners in Syanyonja village do not employ any form of personal protective equipment during mining activities. This could partly be due to lack of sufficient income to purchase them and lack of awareness on the dangers of mercury [21]. In addition, miners are also exposed to other heavy metals such as Copper, Lead and Chromium because these metals are naturally part of auriferous rocks as arseno-pyrite (FeAsS_2), Copper II Sulphide (CuS_2) and Iron (II) Chromium trioxide ($\text{FeO.Cr}_2\text{O}_3$) [38,39]. These metals are released into the tailings during gold processing. Mbonimpa [9] also reported the

presence of heavy metals in samples of rocks and sediments taken from around Namukombe stream and other areas between River Solo and River Malaba in Busia gold district. It is worth noting that most of the respondents as reported previously [13] have explicit fears regarding the health and ecological impacts of mercury usage in gold recovering, peculiarly on the risk of polluting food and water resources and losing fecundity.

3.3 Sources of Water Used for Drinking, Domestic Use and Animal Husbandry

In Syanyonja village, the weather pattern influences a household's choice of the main source of drinking and cooking water. Accordingly, more than half (55%) of the respondents reiterated that they rely on both borehole and Namukombe stream water for drinking and cooking purposes when compared to using borehole water only (Fig. 3). Many residents turn to the stream water when the dry season sets in and the boreholes are dry. Contamination of Namukombe stream, boreholes, springs and bottled water in Uganda are well documented [6,40-45] and thus the results of this study agrees well with studies which indicated that ASGM communities in Uganda have contaminated ecosystems supporting their livelihoods [6,18,46-48]. Water is an indispensable part of life on planet earth and the availability of sufficiently clean and safe water is an important aspect of a community [40].

3.4 Sources of Food

The various sources of food consumed in Syanyonja village are given in Table 3. The major foods consumed are beef, fish, chicken yams, beans, maize and millet flour. A preceding study indicated that fish and yams harvested from Namukombe stream are polluted with mercury [6]. Fish is widely consumed in many parts of Uganda [37,49,50] because it has high protein content, low saturated fat and also contains omega-3 fatty acids, calcium, phosphorus, iron, trace elements like copper and a fair proportion of the B vitamins known to support good health [6,49,51,52]. Besides the good health benefits of fish, there have been many reports on contamination of fish by carcinogenic chemicals in Uganda [6,49,53,54].

Since most of the foods consumed by the community are from within the village, there are high chances that the locals are consuming

mercury-contaminated foods and these risks are expected to be high if most of the fish are consumed whole, especially by children or pregnant women [6]. Other foods such as beef, milk and yams are equally unsafe as they can also get contaminated through water, food and atmospheric deposition of mercury.

3.5 Impacts of ASGM on the environment of Syanyonja village

Field observations and responses gathered reveal that the major environmental problems of ASGM mining in Syanyonja village are deforestation, land degradation, soil erosion, dumping of tailings, air and water pollution (Fig. 2). Deforestation will leave degraded and fragmented habitats for wildlife [55]. More so, this shall increase the vulnerability of forest ecosystems to invasive plant and animal species. Exposure of top soil may initiate soil erosion

which when coupled with the open pits may lead to landslides [55].

Secondary impacts from erosion including sedimentation and siltation are evident in Namukombe stream. Siltation reduces light penetration into water bodies, causing reduced photosynthetic activity in aquatic organisms as well as increased biochemical oxygen demand, clogging of fish gills and thus death of aquatic organisms.

Wildcat pitting in many shallow pits and shafts across the mines were evident and these are excellent breeding grounds for mosquitoes and other disease vectors. In addition, the numerous small shafts and pits dispersed across the mines are hazards for livestock and humans (particularly children), who may fall in the pits, causing loss of lives, income and time. These




Fig. 3. Major water sources used by residents of Syanyonja village

Table 3. Major sources of food for the sampled residents of Syanyonja village

Variable	Categories	Frequency in a month	Percentage
Major sources of proteins	Chicken	02	10
	Fish	10	46
	Beef	12	41
	Others (yams, beans, millet, e.t.c.)	01	3
Sources of fish consumed	Unknown	11	44
	Nearby market	05	20
	Namukombe stream	08	32
	Greater than 10 km from ASGM sites	01	4
Origin of yams consumed	Nearby market	08	32
	Namukombe stream and banks	16	64
	Greater than 10 km from ASGM sites	01	4

observations have been reported in other ASGM sites of Uganda [7,18,48].

It was noted that some parts of Namukombe stream is experiencing siltation due to the cumulative effects of water-intense gold panning. This has greatly increased the turbidity of water from this stream (Fig. 2). Turbidity greater than 5 Formazin Turbidity Unit (FTU) indicate high levels of bacteria, pathogens or particles that can shelter harmful organisms from disinfection [40].

Land is a key denominator in food production and economic development. Artisanal gold mining in Syanyonja is done on land that could otherwise be utilized for agriculture which supports food security. Reluctancy to backfill open pits exacerbate the situation. The interviewed stampeders are incrementally phasing out agriculture and other livelihood enterprises for mining as an economic activity that will salvage them from poverty. It is worth noting that ASGM cause environmental havocs in no small magnitudes [6,16,56] and therefore, as ASGM continue to be a valuable source of income to Ugandans, there is need to blend ASGM with environmental conservation [57].

4. CONCLUSION AND RECOMMENDATIONS

From this survey, the ecological and livelihood challenges associated with ASGM in Syanyonja definitely span beyond the anticipated benefits. Artisanal gold miners are typically poor, hardworking, illiterate with little or no formal education. Miners in Syanyonja are better acquainted with the health impacts of the mercury based ASGM than the accompanying environmental impacts. ASGM in Syanyonja have led to deforestation and pollution of land, air and water. Further research should examine the physicochemical and microbiological profile of the water sources utilized by the residents of Syanyonja village, the mercury levels in the blood, hair and urine of the community as well as the atmospheric flux of mercury in this area. Government should put prohibitive taxes on mercury imports and control its retailing to make mercury unaffordable and unavailable to the artisanal and small-scale gold miners in the move to phase out mercury use in ASGM.

CONSENT

As per international standard, miners' written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

This research required no ethical approval. However, field studies were approved by the Department of Mining and Water Resources Engineering, Faculty of Engineering, Busitema University, Tororo, Uganda (Approval No. BU/UG/2013/95).

ACKNOWLEDGEMENT

Authors TO, KMK & OB are grateful to the World Bank and the Inter-University Council of East Africa (UICEA) for the scholarships awarded to them through the Africa Center of Excellence II in Phytochemicals, Textiles and Renewable Energy (ACE II-PTRE) hosted at Moi University, Kenya.

COMPETING INTERESTS

Authors have declared that no competing interests exist. The products used for this research are commonly and predominantly used products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because the authors do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by any producing company rather it was funded by personal efforts of the authors.

REFERENCES

1. NPA. Second National Development Plan (NDPII). National Planning Authority (NPA), Republic of Uganda. Kampala (UG); 2015.
Available:<http://npa.ug/wp-content/uploads/NDPII-Final.pdf>
2. World Bank. Uganda Overview. Available:<http://www.worldbank.org/en/country/uganda/overview>
3. PwC. Uganda Economic Outlook 2017: Q3; 2017.
Available:<https://www.pwc.com/ug/en/assets/pdf/economic-outlook-2017-third-edition.pdf>
4. Uganda Investment Authority. Mining Sector Profile; 2016.
Available:<http://www.ugandainvest.go.ug/index.php/mining>
5. Omara T, Adupa E, Laker F, Kalukusu R, Otori T. Potential of Sorghum bicolor L. (Moench) and the Effectiveness of Some Organic Amendments in Remediation of

- Petroleum-Oil Vitiated Soils of an Automobile Repair Workshop in Urbanite Kampala. *Asian J Appl Chem Res.* 2019; 3(1):1-10.
Available:<https://doi.org/10.9734/ajacr/2019/v3i130085>
6. Omara T, Karungi S, Kalukusu R, Nakabuye BV, Kagoya S, Musau B. Mercuric Pollution of Surface Water, Superficial sediments, Nile Tilapia (*Oreochromis nilotica* Linnaeus 1758 [Cichlidae]) and Yams (*Dioscorea alata*) in Auriferous Areas of Namukombe Stream, Syanyonja, Busia-Uganda. *PeerJ.* 2019; 7:e7919.
Available:<http://doi.org/10.7717/peerj.7919>
 7. Hinton J, Kabongo I, Kabiswa C, Okedi J & Mbabazi R. The mining and mineralization sector in Karamoja Region: Development opportunities and Constraints. Ecological Christian Organization (ECO), 2011. ISBN 9789970405008
Available:<https://thisafrica.files.wordpress.com/2011/12/mining-sector-report-karamoja-region.pdf>
 8. Nyakecho C, Hagemann SG. An overview of gold systems in Uganda. *Aust J Earth Sci.* 2014;61(1):1-30.
Available:<https://doi.org/10.1080/08120099.2013.831773>
 9. Mbonimpa A. The genesis and characteristics of the gold mineralisation in the area between River Malaba and River Solo in Busia district, South Eastern Uganda. MSc dissertation, Makerere University, Uganda; 2005.
Available:<http://makir.mak.ac.ug/bitstream/handle/10570/2724/Mbonimpa%20Alex.pdf?sequence=1&isAllowed=y>
 10. United Nations Environment Programme (UNEP). Analysis of Formalization Approaches in the artisanal and Small-Scale Gold Mining Sector Based on Experiences in Ecuador, Mongolia, Peru, Tanzania and Uganda. Uganda Case study; 2012.
Available:https://wedocs.unep.org/bitstream/handle/20.500.11822/11357/Formalization_Document_Final_June_2012.pdf?sequence=1&isAllowed=y
 11. Mallongi A, Parkpian P, Pataranawat P, Chinwetkivanich S. Mercury Distribution and its Potential Environmental and Health Risks in Aquatic Habitat at Artisanal Buladu Gold Mine in Gorontalo Province, Indonesia. *Pak J Nutr.* 2015;14(12):1010-2025.
Available:<http://dx.doi.org/10.3923/pjn.2015.1010.1025>
 12. Patricio C, Velasquez-Lopez, Veiga MM & Hall K. Mercury Balance in Amalgamation in Artisanal and Small-scale Gold Mining: Identifying Strategies for Reducing Environmental Pollution in Portovelo-Zaruma, Ecuador. *J Cleaner Prod.* 2009; 18(3):226-232.
Available:<https://doi.org/10.1016/j.jclepro.2009.10.010>
 13. Action Coalition Climate Change (ACCC). Status and Extent of Mercury-use by Artisanal and Small-Scale Gold Miners in Selected Gold Mining Sites of Rubende and Busia Districts, Uganda. Research Paper No. 1, 2018, Kampala, Uganda.
Available:<http://accc-u.org/wp-content/uploads/2018/08/ACCC-REPORT-ON-MERCURY-2018-min.pdf>
 14. Elias C, Deborah SKT, Dewey D, Davey M, Ngallaba SE, Konje E. A Cross-Sectional Survey on Knowledge and Perceptions of Health Risks Associated with Arsenic and Mercury Contamination from Artisanal Gold Mining in Tanzania. *BMC Publ Health.* 2003;13(74):8.
Available:<https://bmcpublihealth.biomedcentral.com/track/pdf/10.1186/1471-2458-13-74>
 15. COWI. Country Reports on Mercury Trade and Use for ASGM. Appendix to the Report "Mercury Trade and Use for ASGM in Sub-Saharan Africa". 2016.
<http://cegemi.com/wp-content/uploads/2016/02/Mercury-Sub-Saharan-Africa-Trade-Report-12-December.pdf>
 16. United Nations Environment Programme. Global Mercury Assessment 2013: Sources, Emissions, Releases and Environmental Transport; UNEP: Geneva, Switzerland; 2013.
Available:<https://www.zaragoza.es/contenidos/medioambiente/onu/942-eng.pdf>
 17. Appleton D, Drasch G, Böese-O'Reilly S, Roider G, Lister R, Taylor H, et al. The GEF/UNDP/UNIDO Global Mercury Project -Environmental and Health Results from a Small-Scale Gold Mining Site in Tanzania; 2005.
Available:<https://core.ac.uk/download/pdf/8793728.pdf>
 18. Ngabiirwe W, Tumwesigye RB, Muloopa HB. Human rights status in the gold supply chain of Uganda: A case for artisanal small-scale mining in Karamoja region. *Global Rights Alert*; 2012.

- Available:https://www.globalrightsalert.org/sites/default/files/Assessment_of_HR_SS-Chain_%20of_%20Minerals_%20A_case_%20ASM_%20in_Karamoja%20.pdf
19. Barreto ML, Schein P, Hinton J and Hruschka F. Understanding the Economic Contribution of small-Scale Mining in East Africa' covering Kenya, Rwanda, and Uganda. Economic Contributions of Artisanal and Small-Scale Mining in Uganda: Gold and Clay; 2018. Available:http://www.responsiblemines.org/wp-content/uploads/2018/03/Uganda_case_study.pdf
 20. Aljazeera. Uganda's gold boom leaves small-scale miners behind; 2019. Available:<https://www.aljazeera.com/ajimpact/uganda-gold-boom-leaves-small-scale-miners-190719030225658.html>
 21. Global Rights Alert. Working conditions of women in mining. A case of Buhweju and Mubende gold mining sites. Available:<https://www.globalrightsalert.org/sites/default/files/The%20Plight%20of%20Women%20in%20Gold%20Mining%20in%20Uganda%20a%20case%20of%20Bubende%20and%20Buhweju%20districts.pdf>
 22. Daily Monitor. Women Exploited in Buhweju Mines; 2018. Available:<https://voice.global/assets/2019/03/The-women-who-toil-in-Buhwejus-gold-pits.pdf>
 23. Minamata Cop 1: Brief on mercury and artisanal and small-scale gold mining (ASGM); 2017. Available:https://ipen.org/sites/default/files/documents/print-EN-ipen-brief-asgm-v1_1-en.pdf
 24. Obase R, Ngoran G, Nde F, Henry L, Ngwane G. Impact of artisanal gold mining on human health and the environment in the Batouri gold district, East Cameroon. Acad J Interdiscipl Stud. 2018;7(1):25-44. Available:<http://doi.org/10.2478/ajis-2018-0003>
 25. World Health Organization. Artisanal and small-scale gold mining and health. Technical paper 1: environmental and occupational health hazards associated with artisanal and small-scale gold mining; 2016. [ISBN 9789241510271] Available:<https://apps.who.int/iris/handle/10665/247195>
 26. Mostafazadeh B, Kiani A, Mohamadi E, Shaki F, Shirazi FH. Mercury exposure of gold mining workers in the northwest of Iran. Pak J Pharm Sci. 2013;26(6): 1267-1270. Available:https://www.researchgate.net/publication/297277587_Mercury_exposure_of_gold_mining_workers_in_the_northwest_of_Iran
 27. Muheki S, Geenen S. Women in (and out of) artisanal mining: a call for revising Uganda's Draft Mining and Minerals Policy. Analysis and Policy Brief no. 25. Institute of Development Policy, University of Antwerp. 2017;6. Available:<https://repository.uantwerpen.be/docman/irua/fb313c/147515.pdf>
 28. Risher JF, Amler SN. Mercury exposure: Evaluation and intervention the inappropriate use of chelating agents in the diagnosis and treatment of putative mercury poisoning. Neurotoxicol. 2005; 26(4):691-699. Available:<https://doi.org/10.1016/j.neuro.2005.05.004>
 29. Grätz T. Moralities, risk and rules in West African artisanal gold mining communities: A case study of Northern Benin. Res Policy. 2009;34(1-2):12-17. Available:<https://doi.org/10.1016/j.resourpol.2008.11.002>
 30. Jaques E, Zida B, Billa M, Greffié C, Thomassin JF. Artisanal and small-scale gold mines in Burkina Faso: Today and Tomorrow. In: Small-Scale Mining, Rural Subsistence and Poverty and West-Africa, Hilson GM ed, Practical Action Publishing: Rugby, UK. 2008;354.
 31. Donkor AK, Bonzongo JC, Nartey VK, Adotey DK. Mercury in Different Environmental Compartments of the Pra River Basin, Ghana. Sci Total Environ. 2006;368(1):164-176. Available:<https://doi.org/10.1016/j.scitotenv.2005.09.046>
 32. Obiri S, Dodoo DK, Essumang DK, Armah FA. Cancer and Non-cancer Risk Assessment from Exposure to Arsenic, Cadmium and Copper by Resident Adults and Children in the Obuasi Municipality, Ghana. Int J Human Ecol Risk Asses. 2010;16(3):651-665. Available:<https://doi.org/10.1080/10807031003788907>
 33. Böse-O'Reilly S, McCarty KM, Steckling N, Lettmeier B. Mercury Exposure and Children's Health. World's Worst Pollution

- Problems Report 2010: Top Six Toxic Threats. Blacksmith Institute/Green Cross, Switzerland. Blacksmith Institute: New York, USA; 2010;201b:33-38, 188-189.
34. Böse-O'Reilly S, Lettmeier B, Matteucci Gothe R, Beinhoff C, Siebert U, Drasch G. Mercury as a serious health hazard for children in gold mining areas. *Environ Res.* 2008;107(1):89-97.
Available: <https://doi.org/10.1016/j.envres.2008.01.009>
 35. Gibb H, O'Leary KG. Mercury exposure and health impacts among individuals in the artisanal and small-scale gold mining community: a comprehensive review. *Environ Health Perspect.* 2014;122(7):667-672.
Available: <https://dx.doi.org/10.1289%2Fehp.1307864>
 36. Ministry of Energy and Mineral Development. Energy and Mineral Development Joint Sector 2012: Ministry of Energy and Mineral Development. Kampala: MEMD. 2012;48-49.
 37. Daily Monitor. Thursday, February 28th, 2019. Miners warned on use of mercury. Available: <https://www.google.com/amp/s/www.monitor.co.ug/News/National/Miners-warned-on-usemercury/688334-5002504-view-asAMP-7gdbdgz/index.htm>
 38. Lacerda LD. Global mercury emissions from gold and silver mining. *Water Air Soil Pollut.* 1997;97:209-221.
Available: <https://link.springer.com/content/pdf/10.1007%2FBF02407459.pdf>
 39. Oyarzun R, Lillo J, Higuera P, Oyarzun J, Maturana H. Strong Arsenic enrichment in sediments from the Elqui watershed, Northern Chile: Industrial (gold mining at El Indio – Tambo district) vs. geologic processes. *J Geochem Expl.* 2004; 84(2): 53-64.
Available: <https://doi.org/10.1016/j.gexplo.2004.03.002>
 40. Omara T, Nassazi W, Adokorach M, Kagoya S. Physicochemical and microbiological quality of springs in Kyambogo University propinquity. *OALibJ.* 2019;6:e5189.
Available: <http://dx.doi.org/10.4236/oalib.1105100>
 41. Haruna R, Ejobi F, Kabagambe EK. The quality of water from protected springs in Katwe and Kisenyi parishes, Kampala city, Uganda. *Afr Health Sci.* 2005; 5:14-20.
Available: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1831893/pdf/AFHS0501-0014.pdf>
 42. Lukubye B, Andama M. Bacterial analysis of selected drinking water sources in Mbarara municipality, Uganda. *J Water Res Protect.* 2017;9: 999-1013.
Available: <https://doi.org/10.4236/jwarp.2017.98066>
 43. Munabi C, Kansime F, Amel A. Variation of water quality in Kakira catchment area, Jinja- Uganda. *Phys Chem Earth Parts A/B/C.* 2009;34:761-766.
Available: <https://doi.org/10.1016/j.pce.2009.06.010>
 44. Kasozi KI, Namubiru S, Kamugisha R, Eze ED, Tayebwa DS, Ssempijja F, et al. Safety of drinking water from primary water sources and implications for the general public in Uganda. *J Environ Public Health.* 2019;12
Article ID 7813962.
Available: <https://doi.org/10.1155/2019/7813962>
 45. Semuyaba AS, Segawa I, Wamala A. Potential risk of lead toxicity from bottled water in Uganda. *Mak Pharm J.* 2014; **12**(1): e13-20.
Available: <https://chs.mak.ac.ug/content/potential-risk-lead-toxicity-bottled-water-uganda>
 46. National Environment Management Authority (NEMA). National State of Environment Report for Uganda: Harnessing our Environment for Wealth Creation, Uganda. 2012;107-108.
 47. New Vision. Museveni stops gold mining in Buhweju; 2013.
Available: https://www.newvision.co.ug/new_vision/news/1327912/museveni-stops-gold-mining-buhweju
 48. NAPE. Small-scale gold miners, government decry poor disposal of mercury in mining.
Available: <http://www.nape.or.ug/news-events/latest-news/45-small-scale-gold-miners-government-decry-poor-disposal-of-mercury-in-mining>
 49. Omara T, Ogwang R, Ndyamuhaki S, Kagoya S, Kigenyi E, Musau B, et al. Spectroscopic analysis of selected priority trace metals in the extant East African gilled lungfish (*Protopterus amphibius*) in Lira municipal lagoon and its edibility health risk. *Sci J Anal Chem.* 2018; 6(5): 38-45.

- Available:<http://doi.org/10.11648/j.sjac.20180605.11>
50. Omara T, Kigenyi E, Laker F, Adokorach M, Otim G, Kalukusu R, et al. Effects of continuous deep-fat frying on the physicochemical properties of assorted brands of edible cooking oils sold in Greater Metropolitan Kampala. *Asian J Appl Chem Res.* 2019;3(2):1-13. Available:<https://doi.org/10.9734/ajacr/2019/v3i230086>
51. Tucker BW. Overview of current seafood nutritional issues: Formation of potentially toxic products, in: Shahidi F, Jones Y, Kitts DD (Eds.), *Seafood safety, processing and biotechnology*. Technomic Publishing Co. Inc., Lancaster, PA (USA); 1997.
52. Omara T. Antibacterial activity of papain hydrolysates of isoelectrically-isolated casein and thermoprecipitated alpha-lactalbumin from bovine and caprine milk on diarrheagenic bacteria. *J Adv Med Life Sci.* 2019;7(3):1-6. Available:<http://doi.org/10.5281/zenodo.2580653>
53. Ogwok P, Muyonga JH, Sserunjogi ML. Pesticide residues and heavy metals in Lake Victoria Nile perch, Lates niloticus, belly flap oil. *Bull Environ Contam Toxicol.* 2009; 82(5):529-533. Available:<http://dx.doi.org/10.1007/s00128-009-9668-x>
54. Birungi Z, Masola B, Zaranyika MF, Naigaga I, Marshall B. Active biomonitoring of trace heavy metals using fish (*Oreochromis niloticus*) as bioindicator species. The case of Nakivubo wetland along Lake Victoria. *Physics and Chemistry of the Earth Parts A/B/C.* 2007;32(15–18):1350–58. Available:<https://doi.org/10.1016/j.pce.2007.07.034>
55. Houdet J, Muloopa H, Ochieng C, Kutegeka S, Nakangu B. *Cost Benefit Analysis of the Mining Sector in Karamoja, Uganda*. Kampala, Uganda: IUCN Uganda Country Office. 2014;9 :82. 978-2-8317-1657-2.
56. Djibril KNG, Tah Bong C, Pierre W, Magha A, Chiane Beng JK, et al. Artisanal gold mining in Batouri area, East Cameroon: Impacts on the mining population and their Environment. *J Geol Mining Res.* 2017;9(1): 1-8. Available:<https://doi.org/10.5897/JGMR2016.0263>
57. Omara T, Othieno N, Obonge J, Ssebulime S, Kansiime M. Characterization and prognostication of wastes generated by industries in Kampala Industrial and Business Park-Namanve. *OALib J.* 2019; 6:e5189. Available:<https://doi.org/10.4236/oalib.1105189>

© 2019 Omara et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://sdiarticle4.com/review-history/52582>